arXiv:1604.04503v1 [math-ph] 15 Apr 2016

A short study of a string on a plane: the energy and the effective mass

S.V. Talalov

Department of Applied Mathematics, State University of Tolyatti,
14 Belorusskaya str., Tolyatti, Samara region, 445020 Russia.
svt_19@mail.ru

Abstract

We investigate the new special class of the finite string on a plane, after the reduction from the relativistic $4D$ case. The suggested special form of the phase space allows to define the extended Galilei group as a group of the space - time symmetry for the considered system. The definition of the energy for the studied non-relativistic string through the Cazimir function of this group is suggested. The concept of the effective mass for the investigated dynamical system is introduced. The appearance of strong correlations between the degrees of freedom even on the classical level is discussed.

keywords: non-critical strings; planar quaziparticles, effective mass
PACS Nos.: 11.25.Pm, 12.90.+b

I. The idea that the point-like singularities of one - dimensional structures in certain crystals can be interpretted as some quasiparticles is very old (see[1], for example). On the other hand, the unconventional points of view on elementary particles continue to appear. So, one of them[2] describes "particles" as defects of the string condensed matter ("string-net condensation"). Speaking about quasiparticles on the plane, we take into account the importance of the "anyon" studies. They are a starting point for the topological quantum calculations. Thus, studies of particle-like excitations of planar infinite - dimensional dynamical systems are of a certain interest. It seems that the string[3] is the simplest mathematical model of this type of system. In the general case, a planar string has cuspidal points that can be interpretted as a certain particle - like structures. There are numerous works devoted to the appearance and the dynamics of these points[4, 5, 6]. However, our approach to the string dynamics is completely original. The main difference is the non-standard hamiltonization of the considered theory. As was noted by Dirac many years ago[7], the various hamiltonian descriptions of one and the same dynamical system lead, in general, to the different physical results. This idea is still actual: the related articles appear from time to time (see[8, 9], for example).

Certain specific models in this aspect have been realised in the earliest author's works[10, 11], where the theory was investigated on both classical and quantum levels. Among other results, it was found that the world - lines of the cuspidal points form the non-trivial braids in the space-time.

In this article we consider the new special class of non-relativistic planar finite strings and suggest a non - canonical approach to the description of the dynamics. As a result, we suggest: firstly, an approach to energy definition and, secondly, the concept of the effective mass for the considered non-relativistic dynamical system. As regards the possible interpretation, we assume that suggested model could be useful for the analysis of some one-dimensional structures on a plane. Examples here could be long polymer molecules, the boundaries between the two continuum media and so on. A separate direction here is the application to development of the string - net hypothesis that was mentioned above (see, also[12]). Indeed, the planar string is the "elementary atom" for this structure.

Let us recall the main points of our approach to the description of the planar string dynamics. The details and proofs can be found in the works cited above. Certain formulae will be modified in accordance with our purposes.

1. Believing in four-dimensionality of the world, we start from the Nambu-Goto string in Minkowski space-time $E_{1,3}$. This object is described by the radius-vector $\mathbf{X} = \mathbf{X}(\xi^0, \xi^1)$ where the dimensionless parameters ξ^0 and ξ^1 describe the evolution and space parametrization of the string. We use the well-known orthonormal gauge so that the equations $\partial_+\partial_-\mathbf{X} = 0$ and the "constraints" $(\partial_\pm\mathbf{X})^2 = 0$ hold. The symbols $\partial_\pm$ denote the derivatives $\partial/\partial\xi_\pm$ where the cone parameters $\xi_\pm = \xi^1 \pm \xi^0$. Apart from the equalities, this point gives our theory the dimensional constant[1] γ and the standard formulae for the dynamical invariants (energy-momentum and angular momenta).

2. The standard boundary conditions for the case $\xi^1 \in [0, L]$ take place:

$$\partial_1\mathbf{X}\Big|_{\xi^1=0} = \partial_1\mathbf{X}\Big|_{\xi^1=L} = 0\,. \tag{1}$$

   The direct consequense of the condition at the point $\xi^1 = 0$ will be the equality $\partial_+\mathbf{X}(\xi) = -\partial_-\mathbf{X}(-\xi)$. Therefore, we can consider the single vector- function on the interval $[-L, L]$

$$\partial\mathbf{X}(\xi) = \varepsilon\partial_\varepsilon\mathbf{X}(\varepsilon\cdot\xi)\,, \tag{2}$$

[1]The proportionality coefficient in the string action.

where $\varepsilon = \varepsilon(\xi) \equiv sign(\xi)$, instead of two functions $\partial_{\pm}\mathbf{X}$ on the interval $[0, L]$. Further we investigate the special case of the finite string so that the parameter $\xi^1 \in [0, \infty)$. The limit $L \to \infty$ conserves the conditions (1), but the value $\lim_{L\to\infty} \partial_0 \mathbf{X}$ is undefined. To ensure the convergence of improper integrals in the subsequent studies, we limit ourselves to the consideration of the special class of string configurations. We suppose that the vector-function $\partial\mathbf{X}(\xi)$ has the form

$$\partial\mathbf{X}(\xi) = \varkappa \exp\Big(-p\,(\xi + q)^2\Big)\mathbf{e}(\xi)\,, \tag{3}$$

where $p > 0$, q and $\varkappa$ are definite constants and the vector function $\mathbf{e}(\xi)$ has bounded components. The zero component of this vector - field must satisfy the condition $\lim_{\xi\to-\infty} e_0(\xi) = 1$. For every value of the parameter ξ the vector $\mathbf{e}(\xi)$ is a dimensionless light - like projective (scale - invariant) vector. Thus the representation (3) separates out the scale-transformed mode $\varkappa$. Speaking about the separation of the scale-transformed mode, this step is quite natural in our opinion because the Nambu-Goto action describes the scale-invariant theory. As regards our subsequent purposes here, we intend to separate the general set of dynamical variables into two groups. The first group (internal variables) will be invariant in relation to both the scale transformations and motions of the target space, the second group (external variables) will be transformed in some way.

3. Let us select an orthonormal basis $\mathbf{e}_\nu(\xi)$ so that $\mathbf{e} = (\mathbf{e_0} - \mathbf{e_3})/2$, $\mathbf{e}_\mu\mathbf{e}_\nu = g_{\mu\nu}$. Obviously, this basis exists but is non - unique. The arbitrariness which arises here will be taken into account later. After that we introduce the vector - matrices $\widehat{\mathbf{E}}$ and $\widehat{\mathbf{E}}_0$ by means of the equalities $\widehat{\mathbf{E}} = \mathbf{e}_\nu(\xi)\mathbf{M}^\nu$ and $\widehat{\mathbf{E}}_0 = \mathbf{b}_\nu\mathbf{M}^\nu$ where the four vectors $\mathbf{b}_\mu$ form a certain stationary orthonormal basis and the four matrices $\mathbf{M}_\mu$ form the "vector" with components $(\mathbf{1}, \boldsymbol{\sigma_i})$. The matrices $\widehat{\mathbf{E}}$ and $\widehat{\mathbf{E}}_0$ are connected by the $SL(2, C)$ - valued field $T(\xi)$ by means of equality

$$\hat{\mathbf{E}}(\xi) = T(\xi)\hat{\mathbf{E}}_\mathbf{0}T^+(\xi)\,.$$

We make the reduction to $D = 1 + 2$ space-time through the requirement $T \in SL(2, R)$. As a consequence, $\mathbf{e}_2(\xi) \equiv \mathbf{b}_2$. Thus the reduced space-time is any space $E_{1,2} \perp \mathbf{b}_2$. In the space $E_{1,2}$, we reconstruct the tangent vectors $\partial_{\pm}\mathbf{X}$ with help of the formulae (2) and (3), where

$$\mathbf{e}(\xi) = \frac{1}{2}\Big[\left(t_{11}^2 + t_{12}^2\right)\,\mathbf{b}_0 - 2\,(\,t_{11}t_{12})\,\mathbf{b}_1 - \left(t_{11}^2 - t_{12}^2\right)\,\mathbf{b}_3\Big]\,. \tag{4}$$

The functions $t_{ij} = t_{ij}(\xi)$ are the elements of the matrix $T(\xi)$. Obviously, this matrix solves the linear problem

$$T'(\xi) + Q(\xi)T(\xi) = 0\,, \tag{5}$$

where the coefficient matrices $Q = -T'T^{-1}$. The arbitrariness that was mentioned above makes it possible to reduce the matrix Q to the form

$$Q(\xi) = -\varrho\,(\xi)\boldsymbol{\sigma}_+ + \varrho\,(\xi)\boldsymbol{\sigma}_-\,.$$

The function $\varrho\,(\xi)$ is real - valued, therefore $T(\xi) \in SO(2)$ after this reduction. The introduction of this function is justified from the geometrical viewpoint: the coefficients of the second fundamental form of the string world - sheet will be proportional to the functions $\varrho\,(\pm\xi_\pm)$. In addition, it must be emphasised that the executed procedure $SL(2,R) \ni T \to T \in SO(2)$ does not commute with the Lorentz boosts in the space - time $E_{1,2}$. Thus, our further theory is non-relativistic. These questions have been investigated in the cited works in detail.

II. Let us fix the type of the functions $\varrho\,(\xi)$. To satisfy our requirement $|\mathbf{e}(\xi)| < const$, we make the following statement:

$$\varrho\,(\xi) = \rho(\xi) + \frac{1}{2}\,\omega\,, \qquad \xi \in (-\infty, \infty)$$

where the real function $\rho(\xi) \in \mathcal{S}$ (Schwarz space) and the real constant ω take arbitrary values. By virtue of these assumptions, the matrix $T(\xi)$ that solves the system (5) has the following form:

$$T(\xi) = \begin{pmatrix} \cos[\,I(\xi) + \omega\xi/2 + \beta\,], & \sin[\,I(\xi) + \omega\xi/2 + \beta\,] \\ -\sin[\,I(\xi) + \omega\xi/2 + \beta\,], & \cos[\,I(\xi) + \omega\xi/2 + \beta\,] \end{pmatrix}, \tag{6}$$

where $I(\xi) = \int_{-\infty}^{\xi} \rho(\eta)d\eta$ and $\beta \in [0, 2\pi)$. Thus, we have a certain set of independent dynamical variables that parametrize the string $\mathbf{X}(\xi^0, \xi^1) = X_1\mathbf{b}_1 + X_3\mathbf{b}_3$ at this (intermediate) stage of our investigations. This set consists of the scale - invariant quantities $\rho(\xi)$, ω, β, p, q and the scale - transformed quantities $\varkappa$ and $\mathbf{Z}$. The latter are the integration constants required for the reconstruction $\partial_\pm\mathbf{X} \to \mathbf{X}$. The group $E(2)$ transforms[2] the variable β and the constant vector $\mathbf{Z}$ only. We emphasize that the identity $\varrho\,(\xi) \equiv 0$ should be fulfilled on any interval $[a, b] \subset (-\infty, \infty)$. This very fact makes it impossible to trivialize the second fundamental form by means of certain conformal transformations $\xi_\pm \to \widetilde{\xi}_\pm = A_\pm(\xi_\pm)$, where $A'_\pm > 0$.

[2]The group of the motions of the plane E_2.

As a result, we have the following expression for the complex - valued function $\mathcal{X} = X_3 + iX_1$:

$$\mathcal{X}(\xi^0, \xi^1) = \mathcal{Z} - \frac{\varkappa}{2} \int\limits_{-\infty}^{\infty} d\eta \Big[\varepsilon(\xi_+ - \eta) - \varepsilon(\xi_- + \eta)\Big] \times$$

$$\times \quad \exp\Big[-p\,(\eta + q)^2 + i\Big(2I(\eta) + \omega\eta + 2\beta\Big)\Big]\,, \qquad \mathcal{Z} = Z_3 + iZ_1\,.$$

Note that the scale - invariant part of the function $\mathcal{X}(\xi^0, \xi^1)$ has the form of the Gabor transform for the function $\Big[\varepsilon(\xi_+ - \eta) - \varepsilon(\xi_- + \eta)\Big] \exp\Big(2iI(\xi)\Big)$. Let us write the Nöether invariants for the considered string:

$$P_\mu = \gamma \int_0^\infty \partial_0 X_\mu \, d\xi^1\,, \qquad M_{\mu\nu} = \gamma \int_0^\infty (\partial_0 X_\mu X_\nu - \partial_0 X_\nu X_\mu)\, d\xi^1\,.$$

Of course, we can use these formulae for the spatial indices only, because our reduction is non - relativistic. The formulae (2), (3), (4) and (6) make it possible to calculate the components P_μ and $M_{\mu\nu}$ through the variables introduced above; in accordance with our assumptions, all integrals converge. The result for the momentum $P_3 + iP_1$, the square of the momentum $\mathbf{P}^2 = P_1^2 + P_3^2$ and the internal angular moment $\mathsf{S} = M_{13} - Z_1 P_3 - Z_3 P_1$ has the form:

$$P_3 + iP_1 = \gamma\varkappa\mathcal{I}_0(\omega)\,, \qquad \mathbf{P}^2 = \gamma^2\varkappa^2\mathfrak{I}_P(\omega)\,, \qquad \mathsf{S} = (\gamma\varkappa^2/2)\mathfrak{I}_S(\omega)\,, \quad (7)$$

where

$$\mathcal{I}_0(\omega) = \frac{1}{2} \int\limits_{-\infty}^{\infty} d\xi \exp\Big[-p\,(\xi + q)^2 + i\Big(2I(\xi) + \omega\xi + 2\beta\Big)\Big]\,,$$

$$g(\omega) \equiv \mathfrak{I}_P(\omega) - i\mathfrak{I}_S(\omega) = \frac{1}{4} \iint\limits_{-\infty}^{\infty} d\xi d\zeta \exp\Big[-p\,\Big((\xi + q)^2 + (\zeta + q)^2\Big)\Big] \times$$

$$\times \quad \left[\cos\left(2\int_\zeta^\xi \rho(\eta)d\eta + \omega(\xi - \zeta)\right) + i\varepsilon(\xi - \zeta)\sin\left(2\int_\zeta^\xi \rho(\eta)d\eta + \omega(\xi - \zeta)\right)\right] \quad (8)$$

To make the formulae shorter, we use the complex - valued functions here. Note also that $\mathfrak{I}_P(\omega) = |\mathcal{I}_0(\omega)|^2 > 0$. Let us exclude the variable $\varkappa$ from the second and third equalities (7). As result we have the equation that connects the quantities $\mathbf{P}^2 = \mathbf{P}^2(\omega)$ and $\mathsf{S} = \mathsf{S}(\omega)$ for the same value of the variable ω:

$$\Phi_1(\omega) \equiv \mathbf{P}^2(\omega)\mathfrak{I}_S(\omega) - 2\gamma\mathsf{S}(\omega)\mathfrak{I}_P(\omega) = 0\,. \quad (9)$$

The "constraint" $\Phi_1(\omega) = 0$ is not the only consequence of the equalities (7). Indeed, if we use the integral representation $\varepsilon(\eta) = (i/\pi)\int_{-\infty}^{\infty}(e^{-i\kappa\eta}/\kappa)\,d\kappa$, we can write the integral formula that connects the quantities $\mathbf{P}^2(\omega)$ and $\mathsf{S}(\omega)$ for the different values of the variable ω:

$$\Phi_2(\omega) \equiv \mathsf{S}(\omega) - \frac{1}{2\pi\gamma}\int\limits_{-\infty}^{\infty}\frac{\mathbf{P}^2(\kappa)}{\kappa-\omega}d\kappa = 0\,. \tag{10}$$

The improper integral is understood here as the Cauchy principal value. To deduce this equation we firstly use

$$\iint_{-\infty}^{\infty} d\xi d\zeta F(\xi,\zeta)\exp\left[-p\left((\xi+q)^2+(\zeta+q)^2\right)\right] = 0$$

for every antisymmetric function $F(\xi,\zeta)$ and, secondly, the identity $(d\varkappa/d\omega) \equiv 0$. Combining equalities (9) and (10) we can easily write the closed singular integral equations for both the internal angular moment S and the square of the momentum $\mathbf{P}^2$. For example, the last value satisfies the equation

$$\mathfrak{I}_S(\omega)\mathbf{P}^2(\omega) = \frac{\mathfrak{I}_P(\omega)}{\pi}\int\limits_{-\infty}^{\infty}\frac{\mathbf{P}^2(\kappa)}{\kappa-\omega}d\kappa\,. \tag{11}$$

The methods for solving such equations are well-investigated [13]. In practice the expression $\mathbf{P}^2 = \gamma^2\varkappa^2\mathfrak{I}_P(\omega)$ gives the special solution of the equation (11) - such solution that depends on the single constant $\varkappa$. Can there be another solution of the equation (11)? As is known, the general solution of this equation contains the number of constants $\varkappa_1,\ldots,\varkappa_n$, where the number $n = \mathrm{ind}\mathsf{G}$ is the index of the function $\mathsf{G}(\omega) = -g(\omega)/g^*(\omega)$ on a closed loop formed by the real ω-axis and the point $\omega = \infty$. Note that the function $g(\omega)$ can be continued analytically to the complex plane for $Im\,\omega > 0$. Because $Re\,g(\omega) > 0$ for the real ω and $g(\infty) = 0$, we have the equalities[13]: $\mathrm{ind}\,g = -\mathrm{ind}\,g^* = 1/2$. Thus $n = 1$ and the considered special solution of the equation (11) is unique. For example, there is no solution of the equation (11) so that $\mathbf{P}^2(\omega) \equiv const$. This very fact is important to us in our further investigations.

III. As noted above, the string energy is not defined for the non-relativistic reduction. In the works[10, 11] the reduction of the space-time $E_{1,2}$ to the space and time $E_2 \times \mathsf{R}$ was supplemented by the consideration of the extended[3] Galilei group instead of the $3D$ Poincare group. After that the

[3]We consider the extension with a single parameter m_0 only. The two - parameter extension of the Galilei group was applied for the anyon modelling in the work[14].

Cazimir function $\widehat{C}_3 = \widehat{H} - (1/2m_0)\widehat{\mathbf{P}}^2$ for the extended Galilei algebra was used to define the energy (the quantities $\widehat{H}$ and $\widehat{\mathbf{P}}$ are the generators of the time and space translations correspondingly). Without going into all details here, we note that the correct definition of the extended Galilei group as a group of the global symmetry was possible because of a special selection of independent dynamical variables. So, the variables P_1, P_3 and S were declared as independent (global) variables instead of the variables $\varkappa$ and β. The equality (9) was considered as a constraint. Recall that the results of the works[10, 11] correspond to the case $\omega = 0$ only. Above we have shown that the dependence of the variables P_1, P_3 and S on the variable ω is not trivial. Therefore, the replacement $(\varkappa, \beta) \to (P_1, P_3; \mathsf{S})$ is possible locally only. To perform this procedure, we consider the phase space $\mathcal{H}$ as the product bundle $\mathcal{H} = \mathcal{H}_2 \times \mathcal{B}$, where the fiber $\mathcal{H}_2$ is the finite - dimensional phase space which corresponds to some structureless free particle on a plane; the fundamental coordinates here are the variables S, $\mathbf{P}$ and $\mathbf{B} = m_0[\mathbf{Z} - (\xi^0/\gamma)\mathbf{P}]$. The base $\mathcal{B} = \mathsf{R}_\omega \times \mathcal{H}_\rho \times \mathcal{H}_{p,q}$ is parametrized by the variables $\omega \in \mathsf{R}_\omega$, $\rho(\xi) \in \mathcal{H}_\rho$ and $(p, q) \in \mathcal{H}_{p,q}$. The extended Galilei group acts independently on each fiber $\mathcal{H}_2$ as a structural bundle group. The equalities (9) and (10) define the indepenent constraints in the global phase space $\mathcal{H}$. From the geometrical viewpoint, the surface W_Φ of these constraints is a non - trivial cross-section of the bundle $\mathcal{H}$. The Poisson brackets are postulated in the form:

$$\{\rho(\xi), \rho(\eta)\} = -\frac{\gamma}{4m_0\mathsf{E}_0}\,\delta'(\xi - \eta)\,, \qquad \{P_i, B_j\} = m_0\delta_{ij}\,, \qquad \{p, q\} = \frac{\gamma}{m_0\mathsf{E}_0}$$

(other possible brackets equal zero). Note that $\{\Phi_1(\omega), \Phi_2(\overline{\omega})\} \equiv 0$. The value E_0 defines the energy scale; together with constants γ and m_0 it is the *in-put* dimensional constant in our theory[4].

Because the Cazimir function C_3 is interpreted as the "internal energy" of a dynamical system, we can write the hamiltonian $H = H(\omega)$ in the form:

$$H(\omega) = \frac{\mathbf{P}^2(\omega)}{2m_0} + h_{int}(\omega) + \sum_i \ell_i\Phi_i(\omega)\,,$$

where the functions ℓ_i, $i = 1, 2$ are Lagrange multipliers. The value $h_{int}(\omega)$ is the energy that corresponds to the "internal" degrees of freedom (p, q), ω and $\rho(\xi)$. It would be natural if we define this quantity as the hamiltonian that generates the dynamics of the corresponding variables. Let the function

[4]Because we reduced our theory from the relativistic case, we can avoid the introduction of the additional constant E_0 assuming that $\mathsf{E}_0 = m_0c^2$. Nevertheless, we conserve the constant E_0 as the independent dimensional constant.

$f(\omega, \mathsf{S})$ be an arbitrary function of the annulators ω and S. Taking into account the definition of the Poisson brackets, the function

$$h_{int}(\omega) \equiv h_{int}[S, \omega;\, p,\, \rho(\xi)] = \mathsf{E}_0 \left(2 \int \rho^2(\xi) d\xi + p \right) + f(\omega, \mathsf{S})$$

provides the dynamics $\rho(\xi^1) \to \rho(\xi^1 + \xi^0)$ and $q \to q + \xi^0$. Indeed, if we define the physical (i.e. dimensional) time t as $t = (m_0/\gamma)\xi^0$, the canonical equations

$$\partial\rho/\partial t = \{h_{int}, \rho\}\,, \qquad \partial q/\partial t = \{h_{int}, q\}$$

take place (for $\ell_1 = 0$). As regards the "observables" – the functions of canonical variables – the dynamics will be correct too. For example, we have the following representation for the angle β:

$$\beta(p, q, \omega, \rho; \mathbf{P}) = \frac{1}{2}\Big(\arg(P_3 + iP_1) - \arg \mathfrak{I}_0[p, q, \omega, \beta; \rho]\Big|_{\beta=0} \Big)\,,$$

so that $\beta \to \beta + \omega\xi^0/2$ and $\partial\beta/\partial t = \{h_{int}, \beta\}$. What is the explicit form for the function $f(\omega, \mathsf{S})$? Let us consider a special configuration when $\rho(\xi) \equiv 0$ and $p \to 0$. This case corresponds to the straight rotating (with the frequency $\Omega = (\gamma/m_0)\omega$) string with the pulsating length, but with the constant moment S. As it seems, the function $f(\omega, \mathsf{S}) = \gamma\omega\mathsf{S}/2m_0$ is adequate for the description of the energy here. Therefore, the final expression for the hamiltonian will be:

$$H(\omega) = \frac{\mathbf{P}^2(\omega)}{2m_0} + \mathsf{E}_0 \left(2 \int_{-\infty}^{\infty} \rho^2(\xi) d\xi + p \right) + \frac{\gamma\omega\mathsf{S}(\omega)}{2m_0} + \sum_i \ell_i \Phi_i(\omega)\,. \qquad (12)$$

As usual, we can substitute the first type constraints after the calculations of all Poisson brackets. Thus, taking into account the formulae (9) and (10), we write the expression for the energy $\mathsf{E}(\omega)$:

$$\mathsf{E}(\omega) = H(\omega)\Big|_{W_\Phi} = \frac{\mathbf{P}^2(\omega)}{2m_{\mathsf{eff}}(\omega)} + \mathsf{E}_0 \left(2 \int_{-\infty}^{\infty} \rho^2(\xi) d\xi + p \right), \qquad (13)$$

where the effective mass m_{eff} is defined as follows:

$$\frac{1}{m_{\mathsf{eff}}(\omega)} \equiv \frac{\partial^2 \mathsf{E}}{\partial |\mathbf{P}|^2} = \frac{1}{m_0} \left[1 + \frac{\omega}{2} \frac{\mathfrak{I}_S(\omega)}{\mathfrak{I}_P(\omega)} \right]\,.$$

Because the inequality $\mathfrak{I}_P(\omega) > 0$ is fulfilled, the value $m_{\mathsf{eff}}(\omega) \neq 0$, $\forall\, \omega \in (-\infty, \infty)$. On the contrary, the case when $m_{\mathsf{eff}}(\omega) \to \infty$ is possible because

the function $\mathfrak{F}(\omega) = \mathfrak{I}_P(\omega) + (\omega/2)\mathfrak{I}_S(\omega)$ can have certain zeros ϖ_m. This fact means that all modes of our dynamical system are strongly correlated when $\omega \to \varpi_m$. The certain properties of the model will be specific near the points ϖ_m. For example, we have the linear dependence $\mathsf{S} = \alpha\mathsf{E} + C$ when $\omega \neq \varpi_m$. But the values S and E become independent when $\omega = \varpi_m$. All these effects require studies on a quantum level, of course. Our experience in such studies[10, 11] gives hope for some interesting results here.

References

[1] Andreev A F *Sov. Phys. Usp.* **19**, issue 2, 137148 (1976);

[2] M. Levin, X.-G. Wen, *Rev. Mod. Phys.* **77**, 871 (2005).

[3] B. Zwiebach, *A First course in String Theory.* (Cambr. Univ. Press, 2004).

[4] Brundobler S., Elser V. *Am Journ Phys.* **60**, 726 - 732 (1992).

[5] Klimenko S. V., Nikitin I. N. *Theor Math Phys.* **114**, 299 - 312 (1998).

[6] Anderson M.R. *The mathematical theory of cosmic strings.* IOP Publishing: Bristol, 2003.

[7] Dirac P.A.M. *Canadian Journ. of Math.* **2**, No 3, 129 (1950).

[8] Magri F. *Journ. Math. Phys.* **19**, No 5, 1156 –1162 (1978); Kulish P.P., Rayman A.G. *Zap. nauch.semin. LOMI* (in Russian), **77**, 134 – 147 (1978).

[9] Marmo G., Scolarici G., Simoni A. and Ventriglia F., *Theor Math Phys.* **144**, 1190 – 1205 (2005).

[10] Talalov S.V. *Int. Journ. Mod. Phys. A* **26**, No 16, 2757 (2011); arXiv: math-ph//1105.0743v1.

[11] Talalov S.V. *The System of Interacting Anyons: A Visual Model Inspired by String Theory.* in book: Progress in String Theory Reseach.Editor: Fred P. Davis, Nova Science Publishers, 2016 (ISBN: 978-1-63484-005-7).

[12] Fidkowski L., Freedman M., Nayak Ch., Walker K., Wang Zh. arXiv: cond-mat/0610583v1 (2006).

[13] Gakhov F.D. *Boundary Value Problems.* Int. Series of Monographs on Pure and Appl. Math., V. 85, Pergamon Press, 1966.

[14] Jackiw R., Nair V.P. *Phys.Lett.* **B480**, 237-238 (2000).